


MODELLING AND IMPLEMENTING TILING RHYTHMIC CANONS IN OPENMUSIC VISUAL PROGRAMMING LANGUAGE

CARLOS AGON
AND
MORENO ANDREATTA

This article presents tools for the construction of Tiling Rhythmic Canons, implemented in the visual programming language OpenMusic. After a brief presentation of OpenMusic, we will focus on the description of the MathTools environment and its most relevant objects which enable a composer to produce several classes of Tiling Rhythmic Canons by means of constraint programming, group factorizations and polynomial representations. Examples of Tiling Rhythmic Canons include canons by translation (from the simplest cases to the Cyclotomic Canons and Vuza Canons) and by augmentation (i.e. canons obtained by affine transformations).

1. The OpenMusic visual programming language and the MathTools environment


OpenMusic (OM) is a visual programming environment dedicated to musical composition, designed and developed at Ircam. It allows for the creation and experimentation with compositional models through programming by means of a graphical interface that represents musical structures and processes. In order to facilitate the understanding of the illustrations in this paper, the basic concept of *patch* will be explained in this section.ⁱ A patch is a graphical representation of a program. In a patch, boxes represent functions, each one having a set of inputs (at the top of the box) and a set of outputs (at the bottom of the box). These boxes can be connected together in order to create the functional layout of the program, i.e. what it will do. OM possesses a number of functions of varying complexity and specialization. New ones may be created by the user. Example 1 shows a simple patch using the functions for the addition and multiplication of integers. It is the graphical equivalent of the expression $(3 + 6) \times 100$.



EXAMPLE 1. A PATCH THAT CORRESPONDS TO THE EXPRESSION $(3 + 6) \times 100$


If the number 3 in the expression $(3 + 6) \times 100$ is made into a variable, the result is a single parameter function defined as follows: $f(x) = (x + 6) \times 100$. Example 2 shows the patch corresponding to the previous function where we have replaced the initial value 3

by a variable x . From a graphical point of view, this variable is represented by an arrow shaped box (at the top of Example 2) whereas the output arrow (at the bottom) specifies the value that results when the patch is calculated.



EXAMPLE 2. A PATCH THAT CORRESPONDS TO THE FUNCTION $f(x) = (x + 6) \times 100$


The patch in Example 2 can in turn be considered as a function call, and can be used in the form of a graphical box in other patches. This makes it possible to create functional abstractions which can then be used in various contexts. Example 3 shows the patch from the previous example in the form of a box, with the number 5 as its argument. The result of the calculation in this patch will therefore be $f(5) / 10 = ((5 + 6) \times 100) / 10 = 110$. This shows the embedding character of the patch object, i.e. the possibility to nest a patch within a second patch (and so on).



EXAMPLE 3. THE RECURSION CHARACTER OF THE PATCH OBJECT


Data structures (classes) can be used in the patches in order to create and manipulate objects (particularly musical ones). These structures are also shown as graphical boxes with inputs and outputs that provide the program access to their content. Example 4 shows a patch for generating a musical sequence algorithmically. It contains a BPF type object (break-point function), a chord sequence (represented in absolute time values)

and a voice (chord sequence in musical rhythmic notation). Among the OpenMusic objects which are most frequently used by composers are the following musical classes: chord, chord-seq, poly (polyphony made up of several voices), midifile, sound (audio file), etc.



EXAMPLE 4. CREATING A MUSICAL SEQUENCE IN A PATCH WHERE THE PITCHES ARE GENERATED FROM THE DATA IN A BREAK-POINT-FUNCTION AND THE RHYTHM IS CREATED OUT OF PROPORTIONAL DURATION RELATIONSHIPS

Each object class possesses a graphical editor, enabling the display and manual editing of musical data. Example 5 shows the graphical editor of the voice type object that was constructed in the previous example.




EXAMPLE 5. A VOICE EDITOR

In addition to standard objects in OpenMusic, we developed a collection of mathematical tools based on algebraic structures, allowing music-theorists and composers to manipulate new musical material, based on the most recent “mathemusal” research.ⁱⁱ These tools are organized in a specialized package called “MathTools”, which associates the previous musical representations with the circular representation, by means of the `n-circle` object.

As firstly observed by a large number of Twentieth-Century music theorists and composers from the American (Milton Babbitt, David Lewin, John Rahn,...) as well as the European (Pierre Barbaud, Iannis Xenakis, Anatol Vieru, Michel Philippot, Guerino Mazzola...) tradition,ⁱⁱⁱ the combinatorial character of the equal-tempered system is formally described by the algebraic model of the cyclic group Z/nZ of order n , where n denotes the number of equal parts in which the octave is divided. It is also common usage, at least in European-oriented computational musicology (Riotte & Mesnage 2006), to make use of the circular representation in order to geometrically capture the algebraic properties of this group within a computer-aided environment. A given chord of m distinct notes (modulo the octave) can therefore be represented from a geometric point of view as a m -polygon inscribed in a circle. To each chord one can associate a sequence of m integer numbers counting the successive intervals in the chord (interval structure). Such a structure is an invariant enabling to identify in a unique way a given chord up to transpositions, as thoroughly studied by Anatol Vieru, whose catalogue of chords is provided by means of this algebraic invariant (Vieru 1980).

Periodic rhythmic patterns can also be represented by means of the `n-circle` object. Example 6 shows two rhythmic patterns which are particularly interesting from a tiling perspective. They are called “3-asymmetric” in Hall and Klingsberger’s terminology (Hall & Klingsberger 2006). From a group-theoretical point of view they are what mathematical music theorists usually call “self-invers partitioning” musical structures. In fact they are “inversionally” related, meaning that they are basically the same orbit under the action of the dihedral group on the twelve-tone system. Moreover, the twelve-tone system can be partitioned into a disjoint union of transposition classes of the two structures. We show their circular representation (by means of the `n-circle` object), the intervallic structure (provided by the `n-structure` function), and their interpretation as chords and one possible rhythmic interpretation via the function `c2rhythm` which maps the circular representation into one corresponding rhythmic pattern (with the choice of a minimal rhythmic value corresponding to the eight note).



EXAMPLE 6. TWO MUSICAL STRUCTURES REPRESENTED AS 4-POLYGONS INSCRIBED IN A CIRCLE AND INTERPRETED AS INVERSIONALLY-RELATED CHORDS AND 3-ASYMMETRIC RHYTHMIC PATTERN

2. A first computational approach to the construction of tiling rhythmic canons: the constraint programming


A rhythmic canon is a polyphony of the same rhythm translated in time. It is defined by two rhythmic patterns: the inner one (R) that is the ground voice (a periodic rhythm) and the outer one (S) made by the entries of each voice. A tiling rhythmic canon is a rhythmic canon where at each pulsation one and only one voice attacks a note. Because of the simplicity of the definition of a Tiling Rhythmic Canon, one might imagine that to modeling this structure as a CSP (Constraint Satisfaction Problem) is a very natural approach. A CSP is given by a triplet (X, D, C) where $X = \{x_1, \dots, x_n\}$ is a finite set of variables, D a set of finite domains, one for each variable (we denote $\delta(x_i)$ the domain of the variable x_i) and C a set of constraints of the form (Z, R) with Z being a subset of $X \{x_1, \dots, x_r\}$ and R a subset of the Cartesian product of their domains, $\delta(x_1) \times \dots \times \delta(x_r) \supseteq R(x_1, \dots, x_r)$. Values $v_{x_1} \in \delta(x_1), \dots, v_{x_r} \in \delta(x_r)$ satisfy a constraint $R(x_1, \dots, x_r)$ if $(v_{x_1}, \dots, v_{x_r}) \in R(x_1, \dots, x_r)$.

An instantiation I is a set of values $(v_{z_1}, \dots, v_{z_r})$ assigned to a subset $Z \subseteq X$. If the inclusion of Z in X is strict, we said that I is a *partial instantiation*. The instantiation I is consistent if for all constraint C_i containing variables in Z the values in I satisfy C_i . A solution is an instantiation consistent with $Z = X$. As an example suppose that we are trying to find a tiling rhythmic canon of 4 voices with period 16. The CSP that model this problem is given by:

- Variables: $R = \{x_1, \dots, x_6\} \cup S = \{y_1, y_2, y_3, y_4\}$
- Domains: $\delta(x_i) = (0 \dots 15) = \delta(y_j)$
- Constraints: $(x_1, \dots, x_6) \otimes (y_1, y_2, y_3, y_4) =$ a permutation of $(0, \dots, 15)$


In order to remove duplicates by translation we add two new constraints, i.e. $x_1 = 0$ and $y_1 = 0$.

Once the CSP is defined, it is given as an input to a constraint solver.^{iv} We obtain as a solution $R = \{0, 1, 4, 5\}$ and $S = \{0, 2, 8, 10\}$ which corresponds to the tiling canon in Example 7. Examples in these papers take the eighth note as the pulsation of the canon.



EXAMPLE 7. A TILING RHYTHMIC CANON OF FOUR VOICES OF PERIODICITY EQUAL TO 16, OBTAINED BY MODELING THE CYCLIC GROUP FACTORIZATION AS A CONSTRAINT SATISFACTION PROBLEM

It is easy to generalize this example as a function taking two parameters p and v specifying respectively the period and the number of voices. An advantage of constraint programming is that it allows searching exhaustively all the solutions for a given period. For instance, for $p = 9$ and $v = 3$ there are 18 canons, as shown in Example 8.




EXAMPLE 8. THE COMPLETE CATALOGUE OF SOLUTIONS OF THE FACTORIZATION PROBLEM OF THE CYCLIC GROUP OF ORDER 9 IN TWO SUBSETS WITH CARDINALITY EQUAL TO 3.

The main problem with constraint programming is the time of computation, which can hinder applications in the factorization of cyclic groups with larger periods (as, for example, 72). It is also difficult to add new constraints to the CSP without changing the performance TIME (as in the family of Vuza or Augmented Canons). For these reasons we need to know more about the problem in order to find more interesting solutions.

3. Tiling Rhythmic Canons and Milton Babbitt's Combinatorality


As already mentioned, the circular representation provides one of the most practical visualization tools for describing the inner symmetries of different kind of musical structures. In our attempt at implementing some computational properties of Set Theory and Transformational Analysis, we initially focused on several families of chord structures having specific symmetries up to transposition, inversion, and complementation (or a combination of the previous properties). These not only correspond to well-known cases in mathematical music theory (Halsey & Hewitt 1978, Vieru 1980) but also have a natural interpretation in terms of tiling rhythmic canons once the pitch content is mapped into the rhythmic domain. As an elementary example let us consider the case of Milton Babbitt's all-combinatorial hexachords. By definition these structures have the property of being at the same time self-complementary (i.e. invariant to complementation) and self-inverse (i.e. invariant to inversion). The OpenMusic patch in Example 9 shows how the six all-combinatorial hexachords are obtained via graphical programming, starting from the two collections of respectively 20 self-inverse and 8 self-complementary chords.



EXAMPLE 9. ENUMERATION AND VISUALISATION OF MILTON BABBITT'S ALL-COMBINATORIAL HEXACHORDS STARTING FROM THE COLLECTION OF THE 20 SELF-INVERS AND THE 8 SELF-COMPLEMENTARY HEXACHORDS


Since each of the six all-combinatorial hexachords tiles the chromatic space by transposition of a given interval (minor second, major second, minor third or triton), it is easy to construct tiling rhythmic canons in two voices having as inner rhythm one of the six all-combinatorial hexachords (mapped in the time domain) and as the outer voice one of the subsets obtained by the previous transposition values. Notice, that among the six previous solutions three are in fact corresponding to rhythmic patterns having a period which is strictly less than 12. In other words, they have transpositional symmetry allowing the reduction of the rhythmic space to be tiled. From the theory of Hajós groups we know that the three non-periodic inner rhythms will generate the tiling in accordance with an outer rhythm corresponding to a periodic subset of $Z/12Z$ of cardinality two. Since the only possibility for such a subset to be periodic is as a subgroup of order two, we deduce that the corresponding tiling canons will have two voices entering regularly with a distance equal to six unities. The patch in Example 10

calculates and visualizes the three tiling rhythmic canons corresponding to the non periodic all-combinatorial hexachords. The function `transp-comb` corresponds to the set-theoretical operation of “transpositional combination” (Cohn 1986) visualizing the tiling process by means of the circular representation.




EXAMPLE 10. AN OPENMUSIC PATCH SHOWING HOW TO OBTAIN THE THREE RHYTHMIC TILING CANONS CORRESPONDING TO NON PERIODIC ALL-COMBINATORIAL HEXACHORDS

The previous example is paradigmatic of a series of examples of tiling rhythmic canons that can be easily calculated since the voices of the canons enter in a regular way, according to the property of one of the factors of the decomposition of Z/nZ being a subgroup of the cyclic group. As we already mentioned, this corresponds to the case of k -asymmetry, a property generalizing Simha Arom and Marc Chemillier’s odditivity (Chemillier 2002). It is easy to show that the traditional definition of a k -asymmetric rhythmic pattern A is equivalent to the condition that $Z/nZ=A+B$, where B is the subgroup of Z/nZ of cardinality equal to k . Example 11 shows how to generate two Tiling Rhythmic Canons starting from the two inversionally-related 3-asymmetric pattern of Example 6.




EXAMPLE 11. TWO TILING RHYTHMIC CANONS BASED ON INVERSIONALLY-RELATED 3-ASYMMETRIC PATTERN

Tiling Rhythmic canons of this type are easy to construct. In fact, one can show that the inner rhythmic pattern of a k -asymmetric rhythm always reduces to the entire cyclic group $\mathbb{Z}/m\mathbb{Z}$ when it is reduced to mod m , where $m = 12/k$. This means that a k -asymmetric rhythm is generated by a permutation of $\{0, 1, \dots, m-1\}$. The following patch (Example 12) shows how to construct all Tiling Rhythmic Canons starting from a k -asymmetric pattern of cardinality equal to 6. In this example $k=3$, which means that the pattern tiles cardinality 6 within the cyclic group $\mathbb{Z}/18\mathbb{Z}$.



EXAMPLE 12. GENERATION OF A TILING RHYTHMIC CANON OF PERIOD 18 STARTING FROM THE 6-ELEMENTS PERMUTATION (0,5,1,3,4,2) INTERPRETED AS A INNER RHYTHMIC PATTERN IN $\mathbb{Z}/18\mathbb{Z}$

Note that by fixing the cardinality m of the inner rhythm (which is equal to 6 in the previous case), the order n of the underlying cyclic group being tiled is always a multiple of m , which immediately provides the information on the number n/m of voices of the canon (hence the order of the asymmetry of the initial pattern). In this way, it is easy to construct canons with big periods, eventually corresponding to one of the values provided by the theory of non-Hajós groups. For example, we can show how to construct a rhythmic tiling canon in 6 voices starting from a 6-asymmetry inner rhythmic pattern, which finally provides a tiling of the non-Hajós group $\mathbb{Z}/72\mathbb{Z}$ (see Example 13). Obviously, this is not a Vuza Canon since one of the factors corresponds to a subgroup of $\mathbb{Z}/72\mathbb{Z}$.



EXAMPLE 13. A TILING RHYTHMIC CANON OF PERIOD 72 CONSTRUCTED FROM THE 12-ELEMENTS PERMUTATION (0,10,1,8,4,2,7,9,5,11,3,6) INTERPRETED AS A INNER RHYTHMIC PATTERN IN $\mathbb{Z}/72\mathbb{Z}$

4. Deriving Tiling Rhythmic Canons from the theory of Cyclotomic Polynomials

The use of polynomials in order to represent the factorization of a cyclic group into subsets goes back to the Minkowski/Hajós problem.^v As originally observed by Rédei (1947), a decomposition of a cyclic group $\mathbb{Z}/n\mathbb{Z}$ into a direct sum $A \oplus B$ of two subsets A and B , can be equivalently expressed in polynomial terms. By putting $A(x) = \sum x^\alpha$, where $\alpha \in A$, the above equation becomes a relation between polynomials with coefficients being either 0 or 1, which will be referred to as “0-1 polynomials”. The factorization condition is expressed by the fact that $A(x) \times B(x) \equiv 1 + x + x^2 + \dots + x^{n-1} \pmod{x^n - 1}$. Factors of the polynomial $\Delta_n(x) = 1 + x + x^2 + \dots + x^{n-1}$, especially those with 0-1 coefficients, are thus of paramount importance which is the main reason to use the theory of cyclotomic polynomials (Amiot *et al.* 2005). More precisely, if $A(x) \times B(x) \equiv 1 + x + x^2 + \dots + x^{n-1} \pmod{x^n - 1}$, then for all $d|n$ (with $d \neq 1$), Φ_d is a divisor of either $A(x)$ or $B(x)$, where Φ_d is the d^{th} cyclotomic polynomial.

In the MathTools environment we have collected a number of tools enabling to obtain Tiling Rhythmic Canons by using cyclotomic polynomials. With exception of some special cases that can be found in the “tiling of the line” literature (and which we discussed in Andreatta *et al.* 2002), it is not known whether a given rhythmic motif enables making a canon unless one is able to exhibit such a canon. Since the establishing of the two Coven & Meyerowitz’ conditions (Coven & Meyerowitz 1998), there is a useful sufficient criterium dealing with the cyclotomic factors. Let $A(x)$ be the associated 0-1 polynomial to an inner rhythm A . We define the two following sets:^{vi}

$$R_A = \{d \in \mathbb{Z}, d > 0, \Phi_d \text{ divides } A(x)\}$$


$$S_A = \{d \in R_A : d = p^\alpha \text{ where } p \text{ is prime and } \alpha \in \mathbb{N}\}.$$

The two Coven-Meyerowitz conditions, called (T1) and (T2), may be expressed in the following way:

(T1) $A(1) = \prod p$ where the product runs over all $p \in S_A$

(T2) If $p^\alpha, q^\beta \dots \in S_A \Rightarrow (p^\alpha \times q^\beta \dots) \in R_A$.


Rhythmic Tiling Canons constructed from a 0-1 polynomial $A(x)$ which has the properties (T1) and (T2) are called “Cyclotomic Canons”. As an example, we show how to construct the same canon of Example 7 with this approach. By choosing as input the value 16, the `cm-conditions` function gives all the possible 0-1 polynomials $A(x)$ associated to subsets of $\mathbb{Z}/16\mathbb{Z}$ having both (T1) and (T2) properties. The sixth solution corresponds to the 0-1 polynomial $A(x) = 1 + x + x^4 + x^5$ obtained by the product of the two cyclotomic polynomials Φ_2 and Φ_8 . The `out-rhythm` function constructs the polynomial $B(x)$ which assures the factorization of $1 + x + \dots + x^{18}$ as $A(x) \times B(x)$ which provides the tiling visualized in the circular representation or as a rhythmic canon, which is the same canon obtained through CSP, as we intended. (Example 14).



EXAMPLE 14. A CYCLOTOMIC CANON OF PERIOD 16 OBTAINED FROM A 0-1 POLYNOMIAL SATISFYING THE TWO COVEN AND MEYEROWITZ CONDITIONS

5. A step-by-step description of Vuza Canon constructions in OpenMusic

The implementation of the algorithm proposed by Vuza in his series of articles published in *Perspectives of New Music* (Vuza 1991-1993), was the first attempt at studying some computational aspects explicitly raised by the author. Before describing the first exhaustive catalogue of solutions that we have obtained in *OpenMusic*, let us shortly provide a general overview of the main tools that we implemented in order to obtain Vuza Canons, shown in Example 15.



EXAMPLE 15. OVERVIEW OF THE MAIN TOOLS IMPLEMENTED IN OPENMUSIC IN ORDER TO OBTAIN VUZA CANONS OF A GIVEN PERIOD

First of all, as we know from the theory of non-Hajós groups, Vuza Canons only exist for some given period n having the property of not being:


- a power p^a of a prime number,
- a product p^2q^2 of the square of two distinct prime numbers
- a product $p \cdot q$ of the power of a prime number by a different prime number
- a product p^2qr of the power of a prime number by two other distinct prime numbers
- a product $pqrs$ of four distinct prime numbers

The function `canon-n` calculates all the periods corresponding to non-Hajós cyclic groups within a given interval (in the previous example, the periods n are constrained between 0 and 300). Every such a period n is decomposable as a product of five numbers $p_1p_2n_1n_2n_3$ where:

- i. p_1 and p_2 are two distinct primes

- ii. $\langle p_1n_1, p_2n_2 \rangle = 1$ (i.e. p_1n_1 is relatively prime with p_2n_2)
- iii. n_1 is an integer greater than 1


The decomposition is obtained by the function `decompo` which takes as input an integer n corresponding to the period of a non-Hajós cyclic group. Note, that the decomposition is not necessarily unique, as the example of 72 shows. Nevertheless, the resulting canons may have the same structural properties in terms of number of voices and number of attacks within each voice (these two values being equal respectively to n_1n_2 and $p_1p_2n_3$). These structural information are provided by the function `infocanon`. By choosing a valid number of voices, the factorization of the group into two non-periodic factors is obtained through the function `patterns` and these sets will finally be provided as input for the function `canons` in order to construct the rhythmic grid of the required Vuza Canon. The representation of the canon in the previous example makes use of the `poly` editor in rhythmic mode. The following example (Example 16) shows two different representations of the same canon by including some information on the pitch content. The first one makes use of the `multi-seq` editor, in which the notes are represented by using proportional durations, whereas the second one uses the same `poly` editor but in a pitch-content mode. In order to show that we are not limited to twelve-tempered equal systems, we map the information of the inner rhythmic pattern into the microtonal space $\mathbb{Z}/72\mathbb{Z}$ corresponding to the 12th of tone division of the octave.^{vii}



EXAMPLE 16. A MICROTUNAL CANON OBTAINED FROM A VUZA CANON OF ORDER 72


6. Canons by augmentation

In this section we will present a generalization leading to canons with augmented voices.^{viii} We will therefore explore a new symmetry of $\mathbb{Z}/n\mathbb{Z}$ under the group of affine transforms. In this new context each symmetry $[a,t](x) := ax + t$ consists of an augmentation with factor a and a translation with summand t . Example 17 shows a rhythmic pattern of period 8 that has been augmented by a factor of 3.



EXAMPLE 17 AUGMENTATION OF A RHYTHM

In order to build an augmented canon, let us shortly provide an overview of the main tools that we implemented, shown in Example 18. Three steps can be identified: In step 1) the user starts by specifying a period p (10, in our example) and the number of attacks compounding the rhythmic pattern (5). The function `ag-canonInfo` provides the list of rhythmic patterns of length 5, together with the multiplicative factors, which need to be applied on them in order to build tiling canons in which each voice is an augmentation of the original pattern according to these multiplicative factors. In Example 18 the solution $((0\ 1\ 3\ 5\ 6)\ ((1\ 3)\ (1\ 7)))$ means that the 5 element rhythmic pattern $(0\ 1\ 3\ 5\ 6)$ can generate a tiling canon both via multiplications by 1 (the original pattern) and by 5 or multiplications by 1 and 7."



EXAMPLE 18. OVERVIEW OF THE MAIN TOOLS IMPLEMENTED IN OPENMUSIC IN ORDER TO OBTAIN AUGMENTED CANONS

At this moment the user must choose the rhythm and the multiplicative factors. Once the choice is made these two parameters are given to the function `allcanons-aff` (step 2). In our example we provide to this function the pattern (0 1 3 5 6) and the factors (1 3). The function `allcanons-aff` provides for the original pattern the list of couples (a_i, b_i) , where a_i are the multiplicative factors and b_i are the translation factors. For example, the solution $((0 1 3 5 6) ((1 0) (3 9)))$ means that the 5 element rhythmic pattern (0 1 3 5 6) can generate a tiling canon via multiplications by 1 and translation by 0 (i.e. the original pattern) and multiplication by 3 and translations by 9 units.

Finally, the function `augmented-canon` uses this information in order to concretely build a rhythmic tiling canon in which all voices are augmentations of the original rhythmic pattern. The period of the augmented canon is given by the period of the original pattern times the least common multiple of the factors ($10 \times 3 = 30$ in Example 18). The augmented voices by 3 are repeated 3 times and translated by 9, 19 and 29.

Note, that for the pattern (0 1 3 5 6) we have another solution with multiplicative factors (1 7), see Example 19. In this case, the result of the function `allcanons-aff` is $((0 1 3 5 6) ((1 0) (7 7)))$ which means that the pattern is completed by 7 voices augmented by a factor 7 and translated by 7, 17, 27, 37, 47, 57 and 67. The period of this canon is $10 \times 7 = 70$.



EXAMPLE 19 AUGMENTED CANON WITH FACTOR 7

7. Conclusions

The problem of constructing tiling rhythmic canons shows the usefulness of a mathematical approach to the formalization of these musical structures. This paper shows in a practical way, how to construct several families of canons in OpenMusic. The reader may consider our references for a more mathematical description of these algorithms.

REFERENCES

Agon, Carlos. 1998. *OpenMusic : Un Langage Visuel pour la Composition Assistée par Ordinateur*, PhD thesis, université de Paris VI.

Agon Carlos, Assayag Gérard, Bresson, Jean. 2006. *The OM Composer's Book 1*, "Musique/Sciences" Series. Sampzon: Ircam/Delatour France.

Assayag Gérard, Rueda Camilo, Laurson Mickael, Agon Carlos and Delerue Olivier. 1999. "Computer Assisted Composition at Ircam: PatchWork & OpenMusic", *Computer Music Journal*, 23(3).

Amiot Emmanuel, Andreatta, Moreno & Agon Carlos. 2005. "Tiling the (musical) line with polynomials: Some theoretical and implementational aspects," *Proceedings of the International Computer Music Conference*, Barcelona.

Andreatta, Moreno. 1999. *La Théorie Mathématique de la musique de Guerino Mazzola et les canons rythmiques*, Mémoire de DEA, Université de Paris IV-EHESS-IRCAM.

Andreatta, Moreno. 2003. *Méthodes algébriques en musique et musicologie du XXe siècle: aspects théoriques, analytiques et compositionnels*, PhD, Ircam-EHESS.

Andreatta, Moreno. 2004. "On group-theoretical methods applied to music: some compositional and implementational aspects," in *Perspectives in Mathematical Music Theory*, (G. Mazzola, E. Puebla et T. Noll 'ed.), EpOs, University of Osnabrück, 122-162.
10

Andreatta, Moreno. 2010. *Mathematica est exercitium musicae. La recherche 'mathémusicale' et ses interactions avec les autres disciplines*. Habilitation Thesis, Institut de Recherche Mathématique Avancée, University of Strasbourg. Available at: <http://recherche.ircam.fr/equipes/repmus/moreno/TexteHDR.pdf>

Andreatta, Moreno, Agon, Carlos & Amiot, Emmanuel. 2002. "Tiling problems in music composition: Theory and Implementation," *Proceedings of the International Computer Music Conference*, Göteborg, 156-163.

Andreatta, Moreno, Agon, Carlos, Noll, Thomas and Amiot Emmanuel. 2006. "Towards Pedagogability of Mathematical Music Theory: Algebraic Models and Tiling Problems in computer-aided composition", *Proceedings of Bridges (Mathematical Connections in Art, Music and Science)*, London, 277-284.

Andreatta, Moreno and Agon, Carlos. 2009. "Guest Editors' Foreword," in *Tiling Problems in Music*, Special Issue of the Journal of Mathematics and Music, Volume 3, Issue 2, 63-70.

Assayag Gérard, Agon Carlos, Laurson Mikael, Rueda Camillo. 1999. "Computer Assisted Composition at Ircam: Patchwork and OpenMusic," *Computer Music Journal*, 23(3).

Bancquart Alain, Andreatta Moreno and Agon Carlos. 2008. "Microtonal Composition," in *The OM Composer's Book 2*, Jean Bresson, Carlos Agon and Gérard Assayag (eds.), "Musique/Sciences" Series. Sampzon: IRCAM/Delatour France, 279-302.

Bloch, Georges. 2006. "Vuza Canons into the Museum," in C. Agon, G. Assayag & J. Bresson (eds), *The OM Composer's Book 1*, Collection "Musique/Sciences". Sampzon: Ircam-Delatour France.

Bresson Jean, Agon Carlos, Assayag Gérard. 2008. *The OM Composer's Book 21*, "Musique/Sciences" Series. Sampzon: Ircam/Delatour France.

Chemillier, Marc. 2002. "Ethnomusicology, Ethnomathematics. The Logic Underlying Orally Transmitted Artistic Practices", in *Mathematics and Music*, G. Assayag, H.G. Feichtinger, and J.F. Rodrigues (Eds), Berlin: Springer Verlag, 2002, pp. 161-183.

Chemillier, Marc. 2007. *Les mathématiques naturelles*. Paris: Odile Jacob.

Cohn, Richard. 1986. *Transpositional combination in twentieth-century music*, PhD thesis, University of Rochester, Eastmann School of Music.

Coven, Ethan and Meyerowitz, Aaron. 1999. "Tiling the integers with translates of one finite set," *J. Algebra*, 212, 161-174.

Fidanza, Giulia. 2007. *Canoni ritmici a mosaico*, tesi di laurea, Università degli Studi di Pisa, Facoltà di SSMMFFNN, Corso di laurea in Matematica.

Friperntinger, Harald. 2001. "Enumeration of non-isomorphic canons," *Tatra Mt. Math. Publ.*, 23, 47-57.

Hall, Rachel and Klinsberg, Paul. 2006. "Asymmetric Rhythms and Tiling Canons," *American Mathematical Monthly*, 113(10), 887-896.

Halsey, David and Hewitt, Edwin. 1978. "Eine gruppentheoretische Methode in der Musik-theorie," *Jahresber. der Dt. Math.-Vereinigung* 80, 151-207.

Noll, Thomas, Andreatta, Moreno, Agon, Carlos, Assayag, Gérard and Vuza, Dan. 2001. "The Geometrical Groove: rhythmic canons between Theory, Implementation and Musical Experiments", *Actes des Journées d'Informatique Musicale*, Bourges, 93-98.

Rédei, Laszlo. 1947. "Zwei Lückensätze über Polynome in endlichen Primkörpern mit Anwendung auf die endlichen Abelschen Gruppen und die Gaussischen Summen," *Acta Math.*, 79, 273-290.

Riotte, André & Mesnage, Marcel. 2006. *Formalismes et modèles musicaux* (in two volumes), "Musique/Sciences" Series. Sampzon: Ircam/Delatour France.

Truchet, Charlotte & Assayag Gérard (eds). 2011. *Constraint Programming in Music*. London: ISTE/Wiley.

Verdi, Luigi. 2008. "The History of Set Theory from a European Point of View," in M. Andreatta, J.-M. Bardez & J. Rahn (eds), *Around Set Theory. A French/American Musicological Meeting*, "Musique/Sciences Series". Sampzon: Ircam/Delatour France.

Vieru, Anatol. 1980. *Cartea modurilor 1*. Bucarest: Ed. Muzicala, 1980. (English translation and enlarged version published as *The book of modes*, Bucarest: Editura Muzicala, 1993).

Vuza, Dan Tudor. 1985. "Sur le rythme périodique," *Revue Roumaine de Linguistique-Cahiers de linguistique Théorique et Appliquée* 23, n°1, 73-103.

Vuza, Dan Tudor. 1988. "Some Mathematical Aspects of David Lewin's Book *Generalized Musical Intervals and Transformations*", *Perspectives of New Music* 26(1), 258-287.

Vuza, Dan Tudor. 1991-1992, "Supplementary Sets and Regular Complementary Unending Canons (in four parts)," *Perspectives of New Music*, 29(2), 22-49.

NOTES

ⁱ For more information on OpenMusic the reader may consult Agon (1998) and Assayag *et al.* (1999). For compositional uses of OpenMusic see the two volumes published in the "Musique/Sciences" Series (Agon *et al.* 2006; Bresson *et al.* 2008).

ⁱⁱ For a short presentation of the "mathemusical" perspective in mathematical music theory see the contribution of one of the authors in the present issue. A more general discussion is contained in Andreatta (2010).

ⁱⁱⁱ For a description of emergence of group-theoretical structures in both European and American traditions, see Andreatta (2003). A historical presentation of classification models for chords within the equal-tempered system in the European perspective is given by Luigi Verdi (2008).

^{iv} See (Truchet and Assayag 2011) for a review of different solvers and their musical applications.

^v See the contribution of one of the authors in the present issue of *Perspectives* for a historical presentation of Minkowski's original Conjecture on the tiling of the Euclidean space by unit cubes and Hajós' solution in terms of group factorizations.

^{vi} In the presentation of Coven & Meyerowitz conditions we follow the simplified notation proposed by Giulia Fidenza (2007). For a more detailed discussion on these two conditions and on the role they are playing with respect to Fuglede's spectral conjecture, see the contribution by Emmanuel Amiot in the present issue of *Perspectives*.

^{vii} For the modelling and implementation of microtonal structures in OpenMusic, see Bancquart (2008).

^{viii} These canons are sometimes referred to as "Noll Canons", by stressing the crucial contribution by Thomas Noll in the development of the theory. Also see Noll *et al.* (2001).